



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 827 743 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
11.03.1998 Bulletin 1998/11

(51) Int. Cl.<sup>6</sup>: **A61K 31/165**, A61K 31/19,  
A61K 38/12

(21) Application number: **97202644.7**

(22) Date of filing: **29.08.1997**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE**

(30) Priority: **04.09.1996 EP 96202460**

(71) Applicant:  
**VRIJE UNIVERSITEIT BRUSSEL  
1050 Brussel (BE)**

(72) Inventors:  
• **Geerts, Albert Emmanuel Corneille**  
1150 Brussel (BE)  
• **Niki, Toshiro**  
1780 Wemmel (BE)

(74) Representative:  
**Eveleens Maarse, Pieter**  
**Arnold & Siedsma,**  
**Advocaten en Octrooigemachtigden,**  
**Sweelinckplein 1**  
**2517 GK Den Haag (NL)**

**1kk1(54)Use of histone decarboxylase inhibitors for treating fibrosis**

(57) The invention relates to a pharmaceutical composition for the treatment of fibrosis.

According to the invention it was surprisingly found that submicromolar concentrations of a histone deacetylase inhibitor and in particular of trichostatin A strongly inhibit three main features of myofibroblastic differentiation of cultured hepatic stellate cells and of skin fibroblasts: (1) synthesis of collagens type I and III, the predominant collagens in fibrosis of liver and other organs; (2) cellular proliferation; and (3) expression of smooth muscle  $\alpha$ -actin, a marker for activated myofibroblasts. These results indicate that trichostatin A is a potent antifibrogenic agent, totally different from any other therapeutic compounds previously described.

**EP 0 827 743 A1**

## Description

The invention relates to a pharmaceutical composition for the treatment of fibrosis, for example liver fibrosis, cirrhosis and fibrotic skin diseases such as hypertrophic scars, keloid and Dupuytren's contracture.

Fibrosis in general is a condition which is characterized by excessive deposition of connective tissue in organs such as skin, lung, heart, liver, pancreas, kidney, etc.

Liver fibrosis is a pathological condition characterized by excessive deposition of connective tissue proteins. Rather than a single disease entity, it is a condition resulting from various diseases, including viral hepatitis, alcoholic liver disease, schistosomiasis, etc. Regardless of the initial cause of the disease, the liver parenchyma is progressively replaced by connective tissue, resulting in deterioration of liver functions.

Skin fibrosis is a pathological condition characterized by excessive deposition of connective tissue proteins in the dermis. Skin fibrosis results of a variety of skin injuries including mechanical trauma, burn wounds, chronic inflammation or chronic auto-immunity. Regardless of the initial cause, the dermis of the skin thickens.

To date, no truly effective therapeutic drug exists for the treatment of fibrotic diseases.

Hepatic stellate cells are the major connective tissue producing cells in both normal and fibrotic livers. In the normal situation, stellate cells serve as vitamin A storage site. These cells are quiescent, show little proliferative activity, and express a limited spectrum of connective tissue proteins. In injured or fibrotic livers, however, stellate cells lose their fat droplets and change their phenotype into myofibroblast-like cells. These myofibroblast-like cells are "activated" cells, show high proliferative activity, and produce large amounts of collagens and other extracellular matrix proteins. Accordingly, this cell type is the logical target for therapeutic intervention. Compounds with antifibrotic effect on stellate cells will be a promising candidate molecule for the treatment of liver fibrosis and cirrhosis.

Fibroblasts of normal skin are quiescent. They synthesize controlled amounts of connective tissue proteins and have low proliferative activity. Following skin injury, these cells become activated, i.e. they proliferate, express  $\alpha$ -smooth muscle actin and synthesize large amounts of connective tissue proteins. The activated cells are often called myofibroblasts.

According to the invention it was surprisingly found that submicromolar concentrations of a histone deacetylase inhibitor and in particular of trichostatin A strongly inhibit three main features of myofibroblastic differentiation of cultured hepatic stellate cells and of skin fibroblasts:

- (1) synthesis of collagens type I and III, the predominant collagens in fibrosis of liver and in many other organs;
- (2) cellular proliferation; and
- (3) expression of smooth muscle  $\alpha$ -actin, a marker for differentiated myofibroblasts. These results indicate that trichostatin A is a potent antifibrogenic agent, totally different from any other therapeutic compounds previously described.

The present invention is directed to the use of a histone deacetylase inhibitor or a pharmaceutical acceptable salt thereof for the preparation of a pharmaceutical composition for the treatment of fibrosis.

Currently six compounds as inhibitors of histone deacetylase are known. Histone deacetylase inhibitors are substances causing an inhibition of the activity of histone deacetylases, resulting in hyperacetylation of histones, comprising:

- trapoxin
- trichostatin A, B, C
- Na-butyrate
- apicidin A (cyclic tetrapeptide)
- HC-toxin (cyclic tetrapeptide)
- chlamydocin.

Sodium butyrate is a natural 4 carbon fatty acid that inhibits histone deacetylase in a non-competitive manner, and requires millimolar concentrations for its biologic activities. Trichostatin A and trapoxin are specific inhibitors of histone deacetylase, and are much more potent than sodium butyrate, being effective in the submicromolar range.

The description in this application is in particular directed to trichostatin A as a non-limiting example and is never intended to limit the scope of the invention.

Trichostatin A or derivatives thereof are disclosed to be useful as an antifibrotic agent for the treatment fibrosis.

Pharmaceutical formulations and use of compounds of Trichostatin A are also disclosed.

Trichostatin A is an antifungal agent originally isolated from *Streptomyces hygroscopicus* by Tsuji et al. (J. Antibiot 29:1-6, 1976). Trichostatin A is also useful as an anticancer (Cancer Res 47:3688-3691, 1987) and an antiprotozoal agent (J. Antibiot 41:461-468, 1988). In the course of experiments we discovered that Trichostatin A has a strong anti-

fibrotic effect on hepatic stellate cells which are the major connective tissue producing cells in the liver.

Trichostatin A can be brought in the form of pharmaceutically acceptable salts. As such pharmaceutically acceptable salts may be used so long as they do not adversely affect the desired pharmacological effects of the compounds. The selection and the production can be made by those skilled in the art. For instance, as a pharmaceutically acceptable salt, and alkali metal salt such as sodium salt or a potassium salt, an alkaline earth metal salt such as calcium salt or a magnesium salt, a salt with an organic base such as an ammonium salt, or a salt with an organic base such as a triethylamine salt or an ethanolamine salt, may be used.

Subjects to be treated by the present invention include both humans and animals.

The antifibrotic agent of the present invention may be administered orally or non-orally. In the case of oral administration, it may be administered in the form of soft and hard capsules, tablets, granules, powders, solutions, suspensions or the like. In the case of non-oral administration, they may be administered in the form of ointments or injection solution, drip infusion formulations, suppositories whereby continuous membrane absorption can be maintained in the form of solid, viscous liquid, or suspension. The selection of the method for preparation of these formulations and the vehicles, disintegrators or suspending agents, can be readily made by those skilled in the art. The antifibrotic agent of the present invention may contain a further substance having antifibrotic activities, in addition to Trichostatin A or its pharmaceutically acceptable salts.

The amount of the active ingredients in the composition of the present invention may vary depending on the formulation, but is usually from 0.1 to 50 % by weight irrespective of the manner of administration. The dose is determined taking into consideration the age, sex, and symptom of the disease of the subject, the desired therapeutic effect, the period for administration, etc. However, preferably a daily dose of the active ingredient is from 0.05 to 100 mg for an adult.

The following examples are provided to illustrate the present invention, and should not be construed as limiting thereof.

Trichostatin A 7-[4-(dimethylamino)phenyl]-N-hydroxy-4,6-dimethyl-7-oxo-2,4-heptadienamide was prepared from the culture broth of *Streptomyces platensis* No.145. Sodium butyrate was purchased from Sigma (St. Louis, MO, USA). Stock solutions of Trichostatin A were prepared in ethanol (2 mg/ml), stored at -20°C, and diluted as required for each experiment. The final concentration of ethanol in the medium was 0.0016%. Stock solutions of sodium butyrate (100 mmol/L) were prepared in distilled water, and diluted as required.

#### Example 1

##### Effect of Trichostatin A and sodium butyrate on the synthesis of collagens type I and III, and smooth muscle $\alpha$ -actin by hepatic stellate cells

The antifibrotic activity of Trichostatin A and sodium butyrate were tested using cultures of hepatic stellate cells.

Stellate cells were isolated from adult Wistar rats (400-550 g) by enzymatic digestion of the liver with collagenase/pronase/DNAase followed by density gradient centrifugation on Nicodenz (Nycomed, Oslo, Norway).

After isolation cells were suspended in Dulbecco's modified Eagle's medium supplemented with 10 % fetal calf serum, 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin, and cultured at 37°C in a humidified atmosphere with 5 % CO<sub>2</sub> and 95 % air.

At day 3 cells were exposed to Trichostatin A (1-100 nmol/L) for 24 h. During the subsequent 24 h cells were metabolically labeled with 25  $\mu$ Ci/ml of Trans <sup>35</sup>S-label (specific activity of <sup>35</sup>S-methionine > 1,000 Ci/mmol, ICN Biomedicals, Costa Mesa CA) while exposure to the compounds was continued. After labelling medium was collected, and subjected to immunoprecipitation using antibodies against collagens type I and III and smooth muscle  $\alpha$ -actin. The precipitates were fractionated by SDS-PAGE and radioactivity of specific bands were measured by the PhosphorImaging technology. For the effect at mRNA level, cells were exposed to 100 nmol/L Trichostatin A for 24 h. RNA was then extracted and analyzed by Northern hybridization analysis.

Table 1 and 2 show the results which were expressed as percentage value relative to the control culture, respectively for Trichostatin A and sodium butyrate. Note the dose-dependent suppression of collagens type I and III synthesis by Trichostatin A. Also, note the strong suppression of smooth muscle  $\alpha$ -actin, an activation marker of stellate cells.

Table 1

Results for Trichostatin A			
	100 nM	10 nM	1 nM
collagen I	37.9±5.6	68.9±4.7	91.7±9.5
collagen III	30.1±9.6	73.2±20.9	71.9±21.0
SM $\alpha$ -actin	15.5±7.4	54.4±5.3	87.6±0.3

Sodium butyrate suppressed smooth muscle  $\alpha$ -actin less effectively, with 50% reduction at a concentration of 1 mmol/L. Inhibition of collagen type III and smooth muscle  $\alpha$ -actin synthesis indicated that Trichostatin A was more potent than butyrate by 5 orders of magnitude.

Table 2

Results for sodium butyrate			
	$10^{-3}$ M	$10^{-4}$ M	$10^{-5}$ M
collagen I	107.8±9.8	127.9±16.2	92.8±11.7
collagen III	67.9±19.1	90.6±42.2	109.4±31.1
SM $\alpha$ -actin	50.0±19.9	91.6±23.4	97.9±24.4

### Example 2

#### Effect of Trichostatin A on the gene expression of collagens type I and III, and smooth muscle $\alpha$ -actin by hepatic stellate cells

Hepatic stellate cells were isolated and cultured as described in Example 1. Cells were exposed to 100 nmol/L Trichostatin A for 24 h. RNA was then extracted and analyzed by Northern hybridization analysis.

At day 3 cells were exposed to 100 nmol/L Trichostatin A or 1 mmol/L sodium butyrate for 24 h. Total RNA was extracted by the method of Chomczynski and Sacchi. Northern hybridization was performed using P-labeled cDNA probes for rat procollagen  $\alpha_1(I)$  (1.6 kb Pst I fragment), rat procollagen  $\alpha_1(III)$  (0.5 kb Hind III/EcoRI fragment), and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (0.5 kb XbaI/HindIII fragment). For smooth muscle  $\alpha$ -actin, a cRNA probe corresponding to the 5'-untranslated region of mouse smooth muscle  $\alpha$ -actin mRNA was used as described previously. The results were quantitated by Phosphor-Imaging and corrected for GAPDH.

The results at the mRNA level are shown in Table 3. Collagen type III and smooth muscle  $\alpha$ -actin mRNA levels were suppressed to a similar extent as at the protein level. On the other hand, only modest tendency for suppression was observed for collagen type I, suggesting that the suppression of collagen type I was mainly posttranslational.

Table 3

	100 nM
collagen I	79.3±13.5
collagen III	39.0±13.5
SM $\alpha$ -actin	20.6±15.4

### Example 3

#### Effect of Trichostatin A and sodium butyrate on the cell proliferation of hepatic stellate cells

Finally, the applicant examined the effects of Trichostatin A and sodium butyrate on proliferation of stellate cells,

since high proliferative activity is one of the major features of myofibroblastic differentiation.

Cells were cultured in triplicate or quadruplicate in 24 well plates (Costar). Cells at day 2 were exposed to 0.01-1 mmol/L sodium butyrate or 1-100 nmol/L Trichostatin A for the 4 subsequent days. Culture medium and test compounds were replaced every day. At day 6 cells were trypsinized and counted in a hemocytometer.

Trichostatin A showed at 100 nmol/L a strong suppressive effect on proliferation. Table 4 summarizes these cell-count results.

Tabel 4

Cell-count results			
control	$10^{-7}$ M Trichostatin A	$10^{-8}$ M Trichostatin A	$10^{-9}$ M Trichostatin A
23.7±1.9	16.2±1.0	22.0±3.0	22.9±1.0
control	$10^{-3}$ M Butyrate	$10^{-4}$ M Butyrate	$10^{-5}$ M Butyrate
23.9±2.0	20.4±0.2	22.2±0.7	21.1±2.0

Finally cells were cultured in triplicate or quadruplicate in 24 well plates. At day 4 cells were exposed tot 0.01-1 mmol/L sodium butyrate or 1-100 nmol/L Trichostatin A for 24 h. Subsequently, medium was changed and cells were further incubated for 20 h with the same concentrations of sodium butyrate or Trichostatin A in the presence of 10  $\mu$ Ci/ml [ $^3$ H]-thymidine (specific activity 25 Ci/mmol, 10  $\mu$ Ci/ml). Radioactivity incorporated into the 2% perchloric acid/95% ethanol/insoluble fraction was measured by scintillation counting. Parallel cultures incubated with [ $^3$ H]-thymidine in the presence of 10 mmol/L hydroxyurea provided the baseline value, which was subtracted from each measurement. Final data were normalized for cell number which was determined by trypsinization of parallel wells. Table 5 shows the results expressed in cpm/cell.

Table 5

[ $^3$ H]-thymidine incorporation			
control	$10^{-7}$ M Trichostatin A	$10^{-8}$ M Trichostatin A	$10^{-9}$ M Trichostatin A
10.0±0.8	1.3±0	9.9±0.4	9.6±0.4
control	$10^{-3}$ M butyrate	$10^{-4}$ M butyrate	$10^{-5}$ M butyrate
9.9±0.4	6.2±0.2	9.5±0.3	9.5±0.2

#### Example 4

##### Effect of Trichostatin A on the cell proliferation of skin fibroblasts

Skin fibroblasts were obtained from male Wistar rats (300-400 g) by the explant tehniue as described previously. All rats were fed ad libitum, and received humane care in compliance with the institution's guidelines for the care and use of laboratory animals in research. Cells were grown in Dulbecco's modified Eagle's medium with 10% fetal calf serum, and cultured at 37°C in a humidified atmosphere with 5% CO<sub>2</sub> and 95% air. When the culture became confluent, cells were trypsinized and replated into 75 cm<sup>2</sup> culture flasks in a split ratio of 1:4. Experiments were performed using confluent cells between passage 5 and 9. Preliminary experiments have shown that under these conditions, skin fibroblasts had acquired the myofibroblast phenotype at the time of experiments, as evidenced by their large size, prominent stress fibers, and expression of smooth muscle  $\alpha$ -actin.

Confluent cultures of skin fibroblasts were exposed to 1, 10 and 100 nmol/L trichostatin A for 24 h. For the TGF- $\beta_1$  experiment, cells were exposed to 5 ng/mL of human recombinant TGF- $\beta_1$  (Calbiochem) and/or 100 nmol/L trichostatin A for 24 h. After the initial 24 h exposure of these compounds, cells were metabolically labeled for 24 h with 50  $\mu$ Ci/mL of Trans  $^{35}$ S-label (specific activity of  $^{35}$ S-methionine>1,000Ci/mmol, ICN Biomedicals, Costa Mesa, CA) in the presence of vitamin C (50 $\mu$ g/mL) (Merck) and  $\beta$ -aminopropionitrile (64  $\mu$ g/mL) (Sigma), while exposure to trichostatin A and/or TGF- $\beta_1$  was continued. Labeled media or cell layers were separately harvested and stored at -70°C. Protein synthesis was measured by trichloroacetic acid (TCA) precipitation. Equal counts (10<sup>6</sup> cpm) of labeled media or cell lysates were subjected to immunoprecipitation. Immunoprecipitation was performed using antibodies against collagen type I

(Southern Biotechnology, Birmingham, AL), type III (Gift by Prof. Dr. D. Schuffan, Freie Univ. Berlin, Deutschland) or smooth muscle  $\alpha$ -actin (clone 1A4, Sigma). After immunoprecipitation, proteins were separated by SDS-PAGE, gels were immersed in Amplify (Amersham, Little Chalfont, UK) and dried, exposed to preflashed autoradiography film (Hyperfilm-MP, Amersham) or quantitatively analyzed by Phosphor-Imaging (Molecular Imager, GS-525, BioRad, USA).

Confluent cultures of fibroblasts were exposed to 100 nmol/L trichostatin A and/or 5 ng/mL TGF- $\beta_1$  for 24 h. Total RNA was extracted by the method of Chomczynski and Sacchi. Northern hybridization was performed as described, using  $^{32}\text{P}$ -labeled cDNA probes for rat procollagen  $\alpha_1(\text{I})$  (1.6 kb Pst I fragment), rat procollagen  $\alpha_1(\text{III})$  (0.5 kb Hind III/EcoRI fragment), and GAPDH (glyceraldehyde-3-phosphate dehydrogenase) (0.5 kb XbaI/HindIII fragment). For smooth muscle  $\alpha$ -actin mRNA was used as described previously. The results were quantitated by Phosphor-Imaging and corrected for GAPDH.

The ratios of protein or mRNA levels of treated vs. controls were calculated for each experimental condition and expressed as means  $\pm$  standard deviation. For the protein study, correction was made for the difference, if any, in the trichloroacetic acid-precipitable counts. The number of experiments used to calculate a mean value was at least 3. The effect was considered statistically significant, when 1.0 did not belong to the 95% confidence interval of the treated/control ratio.

Then we metabolically labeled endogenously produced protein with  $^{35}\text{S}$ -methionine, and immunoprecipitated collagen type I and III using specific antibodies. The immunoprecipitated proteins were separated on SDS-PAGE and quantified by phosphor-imaging. We found that at 100 nmol/L, 10 nmol/L, and 1 nmol/L trichostatin A inhibited synthesis of collagen type I by 30%, 44%, and 17%, respectively (table 6, row 2). Synthesis of collagen type III was inhibited by 41%, 49%, and 10% at the same concentrations of trichostatin A (table 6, row 3). Thus, both collagen type I and III were inhibited by trichostatin A ( $p < 0.05$  for 100 nmol/L and  $p < 0.01$  for 10 nmol/L), with the maximal effect obtained at 10 nmol/L. Synthesis of smooth muscle  $\alpha$ -actin, a marker for myofibroblast differentiation, was also inhibited by  $37 \pm 18\%$ ,  $36 \pm 8\%$ , respectively, following exposure to 100 nmol/L and 10 nmol/L trichostatin A (table 6, row 4). These inhibitory effects of trichostatin A was selective, since over all protein synthesis as measured by TCA-precipitable counts was not affected.

Next we explored whether trichostatin A affected the fibrogenic actions of TGF- $\beta_1$  in skin fibroblasts. For this purpose, cells were exposed to TGF- $\beta_1$  (5 ng/mL) alone, trichostatin A (100 ng/mL) alone, or combination of TGF- $\beta_1$  (5 ng/mL) and trichostatin A (100 nmol/L). Collagen type I and III, and smooth muscle  $\alpha$ -actin were again immunoprecipitated, separated on SDS-PAGE and quantified by phosphor-imaging. As shown in table 7, exposure to TGF- $\beta_1$  stimulated synthesis of collagen type I and III 3.4 and 4.7 fold, respectively. Although stimulation occurred in every preparation of skin fibroblasts, the magnitude of stimulation varied from culture to culture ranging from 1.8 to 5.9 fold for collagen type I and from 2.5 to 8.0 fold for collagen type III. TGF- $\beta_1$  had a modest stimulating effect (1.8 fold increase) on the synthesis of smooth muscle  $\alpha$ -actin. Strikingly, the stimulating effect of TGF- $\beta_1$  was largely abolished, when TGF- $\beta_1$  and trichostatin A were added simultaneously.

Finally, we explored at which level trichostatin A exerted its inhibitory effect on the collagen synthesis by skin fibroblasts. For this purpose we again exposed skin fibroblasts to trichostatin A and/or TGF- $\beta_1$ . After 24 hour exposure, total RNA was extracted and subjected to Northern hybridization analysis. The radioactivity was measured, corrected for GAPDH, and expressed as a ratio to the value of the control culture. As shown in table 8, TGF- $\beta_1$  (5 ng/mL) increased gene expression of collagens type I, and III, and smooth muscle  $\alpha$ -actin 2.3 fold, 2.5 fold and 1.7 fold, respectively. Trichostatin A alone (100 nmol/L) had a modest suppressive effect on the mRNA levels of collagens type I and III and smooth muscle  $\alpha$ -actin. These data suggest that suppressive effect of trichostatin A occurs both at the transcriptional and posttranslational level.

The results of the above mentioned examples are summarized in table 6, 7 and 8.

Tabel 6

The effect of different concentrations of TSA at the protein level. Results are expressed as percentage of control value			
	100 nM	10 nM	1 nM
Collagen I	70 $\pm$ 13%	56 $\pm$ 11%	83 $\pm$ 11%
Collagen III	59 $\pm$ 18%	51 $\pm$ 11%	90 $\pm$ 14%
$\alpha$ -SMA	63 $\pm$ 8%	64 $\pm$ 8%	86 $\pm$ 23 %

Table 7

The effect of TSA on inducing effect of TGF- $\beta$ at the protein level			
	TGF- $\beta$	TGF- $\beta$ +TSA	TSA
Collagen I	347 $\pm$ 185 %	121 $\pm$ 39 %	70 $\pm$ 13 %
Collagen III	476 $\pm$ 273 %	86 $\pm$ 35 %	59 $\pm$ 18 %
SMA	175 $\pm$ 196 %	67 $\pm$ 6 %	63 $\pm$ 8 %

Table 8

The effect of TSA on the inducing effect of TGF- $\beta$ at the mRNA level			
	TGF- $\beta$	TGF- $\beta$ +TSA	TSA
Collagen I	231 $\pm$ 89 %	132 $\pm$ 49 %	84 $\pm$ 14 %
Collagen III	253 $\pm$ 94 %	180 $\pm$ 51 %	74 $\pm$ 10 %
SMA	171 $\pm$ 61 %	114 $\pm$ 24 %	85 $\pm$ 16 %

The results in this application indicate that histone deacetylase inhibitors provide a novel therapeutic potential in the treatment of fibro-proliferative diseases.

In conclusion, we have demonstrated that two unrelated histone deacetylase inhibitors are active antifibronics compounds as two well known experimental models of fibrosis, i.e. hepatic fibrosis and skin fibrosis.

The invention further relates to a method for the treatment of humans or animals afflicted with fibrosis, comprising administering to said subject an effective amount of a histone deacetylase inhibitor in particular Trichostatin A or a pharmaceutical acceptable salt thereof and optionally a suitable excipient.

#### Claims

1. Use of a histone deacetylase inhibitor for the preparation of a pharmaceutical composition for the treatment of fibrosis.
2. Use according to claim 1, wherein the histone deacetylase inhibitor is trichostatin A or a pharmaceutical acceptable salt thereof.
3. Use according to claim 1, wherein the histone deacetylase inhibitor is sodium butyrate.
4. Use according to claim 1-3, wherein the amount of active ingredient varies from 0.1 to 50% by weight.
5. Use according to claim 4, wherein the daily dose of the active ingredient for an adult is from 0.05 to 100 mg.
6. Use according to one of the claims 1-5, wherein the pharmaceutical salt is an alkali metal salt, such as sodium salt or a potassium salt, an alkaline earth metal salt such as calcium salt or a magnesium salt, a salt with an organic base such as an ammonium salt, or a salt with an organic base such as a triethylamine salt or an ethanolamine salt.
7. Method for treating fibrosis comprising administering to humans or animals in need of an anti-fibrotic treatment as anti-fibrotic agent a therapeutically effective amount of a pharmaceutical composition according to claim 1-6.
8. Method for treating fibrosis according to claim 7, wherein the pharmaceutical composition is administered non-orally, in particular in the form of a creme, an ointment, injection solution, drip infusion or suppositories.



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 97 20 2644 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	M. YOSHIDA: "Reversible arrest of proliferation of rat 3Y1 fibroblasts in both the G1 and G2 phases by trichostatin A" EXP. CELL RES., vol. 177, no. 1, 1988, pages 122-131, XP000617221 * the whole document *	1-8	A61K31/165 A61K31/19 A61K38/12
Y	M. YOSHIDA: "Potent and specific inhibition of mammalian histone deacetylase both in vivo and in vitro by trichostatin A" J. BIOL. CHEM., vol. 265, no. 28, 1990, pages 17174-17179, XP000616087 * the whole document *	1-8	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search: see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		13 October 1997	Orviz Diaz, P
CATEGORY OF CITED DOCUMENTS			
<p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &amp;: member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04.007)





European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 97 20 2644

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.8)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	M. KIJIMA: "Trapoxin, an antitumor cyclic tetrapeptide, is an irreversible inhibitor of mammalian histone deacetylase" J. BIOL. CHEM., vol. 268, no. 30, 1993, pages 22429-22435, XP000616088 * the whole document *	1-8	
Y	A.M. GRESSNER: "Effect of n-butyrate on the synthesis of sulfated glycosaminoglycans and hyaluronate by rat liver fat-storing cells (Ito cells)" BIOCHEM. PHARMACOL., vol. 37, no. 19, 1988, pages 3771-3776, XP000645000 * the whole document *	1-8	TECHNICAL FIELDS SEARCHED (Int.Cl.6)



European Patent  
Office

**INCOMPLETE SEARCH  
SHEET C**

Application Number

EP 97 20 2644

Remark : Although claims 7 and 8 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out, based on the alleged effects of the compound/composition.